









Dietary Use of Monoglycerides Blend in Plant-Based Diets Influences Growth Performance, Hepatopancreatic Histology, Immunity and Disease Resistance of Pacific White Shrimp *Penaeus vannamei* Against *Vibrio harveyi*

Romi Novriadi^{1,2*} , Nurul Khotimah¹ , Ahmad Syazni Kamarudin² , Vivi Endar Herawati³ , Slamet Budi Prayitno³ , Seto Windarto³ , Chi Man⁴ , Adriana Barri⁵ , Peer Staehler⁶

¹Department of Aquaculture, Jakarta Technical University of Fisheries (Politeknik Ahli Usaha Perikanan), Ministry of Marine Affairs and Fisheries, Republic of Indonesia, Jl. Raya Pasar Minggu, Jati Padang, Jakarta – 12520, Indonesia

²School of Animal Science, Aquatic Science and Environment, Faculty of Bioresources and Food Industry, Universiti Sultan Zainal Abidin, Aquatic Science and Environment, Terengganu, Malaysia

³Department of Aquaculture, Faculty of Fisheries and Marine Sciences, Diponegoro University. Jl. Prof. H. Soedarto, S.H. Semarang – 50275, Indonesia

⁴BASF East Asia Regional Headquarters Ltd., Hong Kong

⁵BASF, Ludwigshafen, Germany

⁶BASF New Business GmbH, Ludwigshafen, Germany

How to Cite

Novriadi, R., Khotimah, N., Kamarudin, A.S., Herawati V.E., Prayitno, S.B., Windarto, S., Man, C., Barri, A., Staehler, P. (2024). Dietary Use of Monoglycerides Blend in Plant-Based Diets Influences Growth Performance, Hepatopancreatic Histology, Immunity and Disease Resistance of Pacific White Shrimp *Penaeus vannamei* AGAINST *Vibrio harveyi*, 24(6), AQUAST1964. <http://doi.org/10.4194/AQUAST1964>

Article History

Received 26 May 2024

Accepted 19 July 2024

First Online 26 July 2024

Corresponding Author

E-mail: novriadiromi@yahoo.com

Keywords

Monoglycerides

Growth

Innate immune

Penaeus vannamei

Vibrio harveyi

Abstract

Feeding trials were conducted to evaluate the efficacy of commercially available optimized mixtures of monoglyceride blend (MG, as BalanGut® AQ L supplied by BASF, Germany) on growth performance and health condition of Pacific white shrimp *Penaeus vannamei*. A basal diet was formulated without MG, and four other diets were prepared by supplementing the powder forms of MG at inclusion levels of 0.5, 0.75, 1.0, and 1.5%. In a growth trial, eight replicate groups of shrimps were hand-fed with one of the diets four times daily for 60 days and observed for growth, total hemocyte count (THC), lysozyme activity, and histomorphological conditions of the hepatopancreas of the shrimp. In a challenge test, diseases challenged with *Vibrio harveyi* infection at a dose of 1×10^5 CFU mL⁻¹ intramuscularly were conducted and then followed with the observation of the cumulative survival rate after 96 h post-challenge, phenoloxidase, the mean percentage of phagocytic activity and phagocytic index on the blood of the shrimp. In the growth trial, shrimps with inclusion levels of 0.75 up to 1.5% of MG showed significantly higher growth performance than the control treatment. The number of THC and lysozyme activity showed an increasing trend with the inclusion of MG at 0.5 up to 1.0% and then declined with the inclusion of 1.5%. The inclusion of MG did not cause any significant changes in the histomorphology of the shrimp's hepatopancreas. In the challenge test, the lowest survival rate after being infected with *Vibrio harveyi* was found in the control treatment, and the highest survival rate was found in the group of shrimps fed with 1 MG. The phenoloxidase (PO) activity, phagocytic activity, and index were higher in the group of shrimps fed with MG compared to the control treatment. The results of these studies indicate that the dietary supplementation of 0.75 – 1.0 % of MG can improve the growth, health condition, and disease resistance of *P. vannamei*.

Introduction

Pacific white shrimp *Penaeus vannamei* farming industry is challenged by the optimization of feed formulation (Al Eissa et al., 2022; Rahman et al., 2017) and the presence of diseases (Delphino et al., 2022; Geetha et al., 2022) that could affect the sustainability, efficiency, profitability, and productivity of the production system. Feed formula optimization is important since shrimp require specific levels of nutrients for healthy growth at different life stages including during the larvae, post-larvae, juvenile, sub-adults, and adults' stage (Fox et al., 2001). Traditionally, to achieve optimal growth, fish meal (FM) is used to fulfill the specific nutritional requirements of shrimp, which provide a relatively high level of energy per unit weight (Fox et al., 2004; Miles & Chapman, 2006), as an excellent sources of digestible essential amino and fatty acids (Zinn et al., 2009) as well as vitamins and minerals (Nguyen et al., 2009). Due to its balanced nature, FM is the most widely used ingredient in aquafeed formulation, including for shrimp (Abo-Taleb et al., 2021; Tacon, 1993). However, high inclusion of FM is challenging from the perspectives of cost, availability, legislation (Gatlin III et al., 2007) and the sustainability of the environments. Therefore, sustainable feed needs to consider a proper blending of a variety of plant-based ingredients, novel feed ingredients, and animal by-products to substitute the use of FM in the feed formulation (Kok et al., 2020).

Factors to consider in selecting alternative proteins include the quality and economic value of ingredients, sustainability, availability of appropriate nutrients, and attractability or palatability to the shrimp (Novriadi et al., 2019; Rahman et al., 2017). It is also essential to consider limitations on the inclusion level for alternative feed ingredients. A study from Lim and Dominy (1990) showed that as soybean meal (SBM) exceeded 42%, replacing FM in the diet formulation negatively affected the weight gain, feed intake and body phosphorus percentage of the shrimp. An additional study with other alternative ingredients, such as dietary dried distiller's grains with solubles (DDGS), demonstrated that the inclusion of DDGS up to 12% to replace dietary FM also caused adverse effects on the growth performance of shrimp *P. vannamei* (Gyan et al., 2021). These studies indicate that alternative proteins' direct replacement of FM can be challenging. However, proper combination with other ingredients supplemented with additives might provide beneficial impacts to reduce the detrimental effects (Novriadi et al., 2022; Xie et al., 2018)

Other than growth performance, the use of alternative ingredients in combination with novel and functional ingredients can be designed to stimulate and enhance the non-specific immune system of shrimp (Ayiku et al., 2020; Novriadi et al., 2023). The presence of diseases due to parasites, bacteria and viruses continues to cause substantial economic losses worldwide (Jithendran et al., 2021; López-Téllez et al.,

2020; Novriadi, 2016; Shinn et al., 2018). The inclusion of several functional ingredients or additives in the diet formulation could become an essential mitigation approach against pathogens, including the use of yeast (Jin et al., 2018), hydrolyzable tannins (Novriadi et al., 2021), algae (Niu et al., 2019) and bile acids (Niu et al., 2019). However, only a few studies have looked into the evaluation on the efficacy of optimized mixtures of monobutyryl, glycerides, and glycerol as a functional ingredient to support the growth and the non-specific immune system response of shrimp *P. vannamei* (Bruno Corrêa da Silva et al., 2016; Bruno Correa da Silva et al., 2016). Therefore, this study was conducted to investigate the effects dietary blend of commercial monoglycerides (MG, as BalanGut® AQ L supplied by BASF, Germany) supplemented into shrimp feed formulation on the growth performance, feed conversion ratio (FCR), total hemocyte count (THC), lysozyme activity and histomorphological condition of the hepatopancreas. In addition, a challenge test was performed to evaluate the efficacy of MG to support the optimal survival rate of shrimp against infection with *Vibrio harveyi* for 96 hours, followed by the analysis of phenoloxidase activity, phagocytic activity and phagocytic index of the Pacific white shrimp *P. vannamei* at day-5 post-infection.

Materials and Methods

Experimental Diets

Five experimental diets for growth trial and challenge test were formulated according to a typical commercial formulation for Pacific white shrimp *Penaeus vannamei* in Indonesia. The control diet was formulated to contain a mixture of 41% soybean meal (SBM), 12% Poultry meal (PM) and 10% fish meal (FM) as the sources of protein in the diet formulation without monoglycerides (MG). Four experimental diets were formulated to contain increasing levels of MG (0.5%; 0.75%; 1%; and 1.5%) to partially replaced the wheat flour as the filler in diet formulation, and labeled with 0.5; 0.75; 1.0; and 1.5 MG, respectively (Table 1). All ingredients were grounded and passed through < 200 mesh sieve (Jinan Shengrun, China). All dry ingredients were carefully weighed and mixed in a paddle mixer (Marion Mixers, Inc., Marion, IA, USA) in a 100 kg batch, followed by grinding to a particle size of <200 µm using a disk mill (Jinan Shengrun China). Fish oil was then gradually added and mixed constantly. A twin extruder (Jinan shengrun, China) was used to extrude the feed through a 2 mm die at temperature gradient of 62°C, 80°C and 110 °C in three zones of extruder barrel and the die head, respectively. Diets were air-dried in a pulse bed dryer (Jinan Shengrun, China) and stored at 4°C in sealed containers until further use. Proximate and amino acid profiles of the experimental diets were analyzed at Saraswanti Indo Genetech Laboratory, Bogor, West Java, Indonesia and summarized in Table 2.

Growth Trials

Growth trials were carried out at the Faculty of Fisheries and Marine Science, Diponegoro University (Semarang, Central Java, Indonesia). The juvenile stage of the shrimp was tested free from pathogens prior to transport and was nursed in the faculty facilities by utilizing 2000-L tanks with seawater under controlled conditions (Temperature 28 – 29°C, salinity 25 ppt and dissolved oxygen above 4 mg L⁻¹). At the start of the trial, Shrimp (1.06±0.01 g initial mean weight) were randomly distributed into 40 aquaria tank (70 x 35 x 40 cm per aquaria tank). Eight replicate groups of shrimps per dietary treatment were administered different types of experimental diets for 60 days and fed by hand four times daily at 07:00, 11:00, 15:00, and 20:00h. Based on our historic results, feed inputs were pre-programmed assuming the normal growth of shrimp and feed conversion ratio of 1.5. Daily allowances of feed were adjusted based on observed feed consumption, weekly shrimp counts, and mortality. During the 60 days of the feeding trial, water quality parameters, including pH, dissolved oxygen (DO), water temperature, total dissolved solid and salinity were measured four times daily using sensors (Aqua TROLL 500 Multiparameter Sonde instrument) and the data were stored to an application (AquaEasy apps, Bosch, Singapore) for data traceability and recording. Meanwhile, total ammonia-nitrogen (TAN), alkalinity, nitrate, nitrite, ammonia and phosphate were measured once a week using absorption spectrophotometry (DR890, HACH,

USA). At the termination of the feeding period, shrimp in each aquaria tank were group-counted and weighed to calculate the final biomass, final weight, percentage weight gain (PWG), feed conversion ratio (FCR), percentage survival (SR), and percentage (%) thermal unit growth coefficient (TGC) as follows:

$$PWG = \frac{(\text{average individual final weight} - \text{average individual initial weight})}{(\text{average individual initial weight})} \times 100$$

$$FCR = \frac{\text{feed given (g)}}{\text{alive weight gain (g)}}$$

$$SR = \frac{\text{final number of shrimp}}{\text{initial number of shrimp}} \times 100$$

$$TGC = \frac{FBW^{1/3} - IBW^{1/3}}{\Sigma TD} \times 100$$

Where FBW is final body weight, IBW is initial body weight, T is water temperature (°C) and D is number of trial days

Total Haemocyte Count

At the end of the growth trial, hemolymph was sampled from one shrimp per aquaria tank or eight shrimp per dietary treatment to determine the total hemocyte count. Hemolymph (100 µL) of individual shrimps was withdrawn from the pleopod base of the second abdominal segment with a sterile 1-mL syringe (25 G x 13 mm needle). Before hemolymph extraction, the syringe was loaded with a precooled (4°C) solution (10%-EDTA, Na) used as an anticoagulant. The

Table 1. Ingredient composition (% *as is*) of experimental diets containing several inclusion levels of monoglycerides mixture (BalanGut®, BASF, Germany)

Ingredient	Control	0.5 MG	0.75 MG	1.0 MG	1.5 MG
Soybean meal ¹	41.00	41.00	41.00	41.00	41.00
Poultry Meal (pet food grade) ²	12.00	12.00	12.00	12.00	12.00
Monoglyceride³	0.00	0.50	0.75	1.00	1.50
Fish Meal Chile ²	10.00	10.0	10.0	10.0	10.0
Soya Lecithin ⁴	1.00	1.00	1.00	1.00	1.00
Fish Oil ²	3.20	3.20	3.20	3.20	3.20
MCP ²	0.80	0.80	0.80	0.80	0.80
Cholesterol ⁴	0.01	0.01	0.01	0.01	0.01
Wheat flour ⁵	29.79	29.29	29.04	28.79	28.29
L-Lysine HCl ⁴	0.25	0.25	0.25	0.25	0.25
DL-Methionine ⁴	0.35	0.35	0.35	0.35	0.35
L-Threonine ⁴	0.18	0.18	0.18	0.18	0.18
Mineral premix ⁶	0.50	0.50	0.50	0.50	0.50
Vitamin premix ⁷	0.50	0.50	0.50	0.50	0.50
Choline Chloride ⁴	0.20	0.20	0.20	0.20	0.20
Anti-Mold ⁴	0.12	0.12	0.12	0.12	0.12
Stay C – 35 ⁴	0.10	0.10	0.10	0.10	0.10
Sum	100.00	100.00	100.00	100.00	100.00

¹ De-hulled solvent extract soybean meal, PT FKS Multi Agro, Tbk. Jakarta, Indonesia ²

² PT FKS Multi Agro, Tbk. Jakarta, Indonesia

³ BalanGut, A Q L, BASF, Germany

⁴ PT Fianza Putra Perkasa, Jakarta, Indonesia

⁵ PT Pundi Kencana, Cilegon, Banten, Indonesia

⁶ Trace mineral premix (g/100 g premix): cobalt chloride, 0.004; cupric sulfate pentahydrate, 0.550; ferrous sulfate, 2.000; magnesium sulfate anhydrous, 13.862; manganese sulfate monohydrate, 0.650; potassium iodide, 0.067; sodium selenite, 0.010; zinc sulfate heptahydrate, 13.193; alpha-cellulose, 69.664.

⁷ Vitamin premix (g/kg premix): thiamin-HCl, 4.95; riboflavin, 3.83; pyridoxine-HCl, 4.00; Ca-pantothenate, 10.00; nicotinic acid, 10.00; biotin, 0.50; folic acid, 4.00; cyanocobalamin, 0.05; inositol, 25.00; vitamin A acetate (500,000 IU/g), 0.32; vitamin D3 (1,000,000 IU/g), 80.00; menadione, 0.50; alpha-cellulose, 856.81

hemolymph with an anticoagulant solution was diluted in 150 µL of formaldehyde (4%), and then 20 µL was placed on a hemocytometer (Neubauer) to determine THC using an optical microscope (Olympus, DP72).

Lysozyme Activity Analysis

Eight shrimps per dietary treatment were collected at the end of the growth trials to analyze the lysozyme activity levels after 60 days of the feeding period. Commercial kits (Sigma-Aldrich, Cat. No. LY0100) were used to measure the activity in shrimp, and the determination procedures came from the instruction manual. The lysis of the *Micrococcus lysodeikticus* cells defined the results of lysozyme activity. The reactions were conducted at room temperature, and the absorbance of samples at 450 nm was measured using an ultraviolet/visible spectrophotometer (Perkin Elmer, Lambda XLS, USA)

$$\text{Lysozyme activity} \left(\frac{\text{Units}}{\text{mL}} \right) = \frac{(\Delta A_{450} / \text{min Test} - \Delta A_{450} / \text{min Blank}) (df)}{(0.001)(0.03)}$$

df = dilution factor

0.001 = ΔA₄₅₀ as per the unit definition

0.03 = Volume (in mL) of enzyme solution

Histomorphological Condition of Hepatopancreas

Upon termination of growth trials, the hepatopancreas sections from four shrimps per dietary treatment were immediately preserved in Davison’s fixative solution for 48 h at room temperature (Bell and Lightner, 1988) and then transferred to 50% ethanol

solution (VWR, Radnor, PA, USA) until processed by standard histological analysis procedures. Samples were dehydrated through a standard ethanol series to 100%, embedded in paraffin wax, and sectioned at 4 µm intervals for staining with Hematoxylin-Eosin (H&E) stain (Merck, Darmstadt, Germany). Double-blinded evaluation with a grading scale of 1 to 5 was used for estimations. Score 1 was considered the normal condition; subsequent scores accounted for increasing levels of histopathological alteration compared to the normal condition. Images were acquired using a digital imaging microscope (Olympus BX41, Olympus Optical Co., Ltd., Tokyo, Japan).

Challenge Test

In parallel with the growth trial, a separate study explored the resistance of *P. vannamei* fed with MG against pathogenic strain of *Vibrio harveyi* ATCC 14126 and the immune system characteristics post-challenge. For this purpose, 80 shrimp per dietary treatment (1.06±0.01 g initial mean weight) were challenged by intramuscular injection with an LD50 dose of 10⁵ CFU of *V. harveyi* shrimp⁻¹ after receiving the experimental diet for 30 days. The concentration was based on previous work from Novriadi et al. (2022) and shrimp mortality was monitored every 12 – 96 h for a period of a maximum of 5 days. The protective effects of bacteria were evaluated based on the following relative percentage survival (RPS) value (Amend, 1981). In this challenge test, the group of shrimps fed with a control diet without challenge were also included as the negative control group

Table 2. Detail composition of proximate and amino acid (AA) (% as is, dry matter basis) of experimental diets utilized in this trial

	Diet code					Methods
	Control	0,5 MG	0,75 MG	1 MG	1,5 MG	
Proximate analyses (% as is)*						
Ash content	10.68	9.58	9.73	9.92	9.93	Incineration
Fat content	6.68	6.71	6.74	6.75	6.68	Soxhlet extraction
Moisture content	7.43	7.88	7.50	7.18	7.95	Drying method
Protein content	37.76	37.78	37.75	37.84	37.85	Kjeldahl method
Amino acid profile (% as is)*						
L-Serine	1.45	1.49	1.46	1.55	1.49	UPLC Photodiode Array (PDA)
L-Glutamic acid	4.45	4.45	3.36	4.47	4.44	UPLC Photodiode Array (PDA)
L-Phenylalanine	1.75	1.73	1.20	1.88	1.82	UPLC Photodiode Array (PDA)
L-Isoleusine	1.46	1.63	0.99	1.54	1.47	UPLC Photodiode Array (PDA)
L-Valine	1.64	1.65	1.23	1.67	1.73	UPLC Photodiode Array (PDA)
L-Alanine	2.26	2.29	1.28	2.28	2.27	UPLC Photodiode Array (PDA)
L-Arginine	2.76	2.89	1.71	2.83	2.76	UPLC Photodiode Array (PDA)
Glycine	2.80	2.73	1.75	2.80	2.79	UPLC Photodiode Array (PDA)
L-Lysine	1.57	1.64	1.59	1.59	1.59	UPLC Photodiode Array (PDA)
L-Aspartic Acid	2.93	2.94	2.94	2.94	2.89	UPLC Photodiode Array (PDA)
L-Leusine	2.65	2.68	2.67	2.86	2.80	UPLC Photodiode Array (PDA)
L-Tyrosine	1.27	1.34	1.38	1.76	1.27	UPLC Photodiode Array (PDA)
L-Proline	1.99	2.02	1.99	1.97	1.99	UPLC Photodiode Array (PDA)
L-Threonine	1.67	1.69	1.74	1.72	1.73	UPLC Photodiode Array (PDA)
L-Histidine	0.73	0.82	0.82	0.75	0.75	UPLC Photodiode Array (PDA)
L-Tryptophan	0.20	0.26	0.19	0.22	0.20	UPLC Photodiode Array (PDA)
L-Systine	2.14	2.17	2.42	2.33	2.44	UPLC Photodiode Array (PDA)
L-Methionine	1.09	1.13	1.12	1.09	1.12	UPLC Photodiode Array (PDA)

*Analysis conducted by the Saraswanti Indo Genetech Laboratory, Bogor, West Java, Indonesia. Website www.siglaboratory.com

Phenoloxidase Activity, Phagocytic Activity and Index

Upon termination of the challenge test, the phenoloxidase (PO) activity of shrimp was measured spectrophotometrically, recording the formation of dopachrome produced from L-dihydroxyphenylalanine (L-DOPA) (Huang et al., 2010). For this purpose, A total of 100 μL of hemolymph was centrifuged at 700 g (4°C) for 20 min. The supernatant was removed, while the remaining pellets was diluted in 100 μL of cacodylate citrate buffer. A total of 100 μL of the mixture was transferred to a 96 wells-microplate, and added with 50 μL Trypsin (Sigma Aldrich), incubated at room temperature for 10 min, and then added with a 50 μL of L-DOPA. PO activities measurement was made using a microplate reader (R-Biopharm Well Reader, Germany) at 490 nm

Phagocytic activity (PA) and index (PI) were measured using ten shrimp per dietary treatment following the protocols described by Romi (2023). The PA and PI evaluation were described as follows:

$$PA (\%) = \frac{\text{number of engulfing phagocytes}}{\text{number of observed phagocytes}} \times 100\%$$

$$PI (\%) = \frac{\text{number of engulfing particles}}{\text{number of engulfing phagocytes}} \times 100\%$$

Statistical Analysis

Data for growth parameters, total hemocyte counts, lysozyme activity, challenge test, phenoloxidase activity, and phagocytic activity and index were analyzed using regression and one-way analysis of variance (ANOVA) to determine significant differences among treatments followed by Tukey's multiple comparison tests to determine the difference between treatment means among the treatments. Score data for the histomorphological condition of the hepatopancreas of shrimp were treated as categorical data, tested for normality and homoscedasticity, and subsequently analyzed using a linear regression model. All statistical analyses were conducted using the SAS system (V9.4. SAS Institute, Cary, NC, USA).

Results

Water Quality Parameters

Data for water quality analysis were presented in Table 3. During the growth trial, water quality parameters including pH, dissolved oxygen (DO), temperature and salinity were in the range of 7.71 ± 0.18 ; $5.76 \pm 0.24 \text{ mg L}^{-1}$; $27.78 \pm 0.28^\circ\text{C}$; and $27.99 \pm 0.18\%$, respectively. Additionally, ammonia ($\text{NH}_3\text{-N}$), nitrite ($\text{NO}_2\text{-N}$), nitrate ($\text{NO}_3\text{-N}$) and phosphate ($\text{PO}_4\text{-N}$) were $0.02 \pm 0.04 \text{ mg L}^{-1}$; $0.011 \pm 0.005 \text{ mg L}^{-1}$; $4.077 \pm 0.622 \text{ mg L}^{-1}$; and $0.011 \pm 0.005 \text{ mg L}^{-1}$, respectively. Overall, all water quality remained within the acceptable ranges for shrimp growth under outdoor pond conditions.

Growth Performance

Table 4 shows the growth performance of shrimp fed with graded levels of MG. After 60 days of the feeding trial, final biomass (FB), final mean weight (FMW), weight gain (WG) and feed conversion ratio (FCR) of shrimp fed with 0.75%, 1% and 1.5% MG were significantly improved compared to the control treatment ($P < 0.05$). Meanwhile, no significant differences were observed on survival and thermal growth coefficient (TGC) of shrimp fed with the experimental diets.

Total Haemocyte Counts (THC) and Lysozyme Activity

Data for THC and lysozyme activity are presented in Figure 1 and 2. Numerically, THC and lysozyme activity were higher in the group of shrimps fed with 0.75 and 1.0 of MG. However, there were a decreasing trend as the inclusion of MG increased in the diet formulation.

Histomorphological Condition of Hepatopancreas of Shrimp

Figure 3 shows the hepatopancreas of the shrimp *P. vannamei* after feeding the shrimp with experimental diets for 60-days. The results of the histopathological

Table 3. Water quality data during the controlled feeding trials for 60 days in aquaria tank. Data were presented as mean \pm standard deviation (range)

Parameters	Unit	Analysis results
Parameters	Unit	Results
Dissolved oxygen	mg L^{-1}	5.76 ± 0.24
Temperature	$^\circ\text{C}$	27.78 ± 0.28
Salinity	$\%$	27.99 ± 0.18
pH		7.71 ± 0.18
Ammonia ($\text{NH}_3\text{-N}$)	mg L^{-1}	0.02 ± 0.04
Nitrite ($\text{NO}_2\text{-N}$)	mg L^{-1}	0.011 ± 0.005
Nitrate ($\text{NO}_3\text{-N}$)	mg L^{-1}	4.077 ± 0.622
Posphate ($\text{PO}_4\text{-N}$)	mg L^{-1}	0.011 ± 0.005

analysis showed that the addition of MG in the range of 0.5 – 1.0 % MG had a normal structure in the presence of B, R, and F cells compared to control and 1.5% MG. In all dietary treatments, numbers of vacuoles appeared in the tubular epithelial cells of the hepatopancreas of the shrimp.

Challenge Test

After 24 h infection by *Vibrio harveyi*, the tested shrimp showed apathetic behavior near the culture tank, slow movement, experienced damage in the antennae and rostrum. At the end of the challenge test, the cumulative survival rate of shrimp fed with MG was significantly higher compared to the control treatment (Figures 4A and 4B).

Phenoloxidase (PO) Activity and Analysis of Phagocytic Activity and Index

Phenoloxidase (PO) activity at day-5 post injection with *Vibrio harveyi* were displayed in Figure 5. The PO activity was higher in all the group of shrimps fed with MG compared to the control treatment ($P < 0.05$). The mean percentage of phagocytic activity (PA) and phagocytic index (PI) of post challenged shrimp can be viewed in Figure 6. Shrimp fed diets containing MG, at

any level, had significantly ($P < 0.01$) higher phagocytic activity compared to the control treatment. In detail for PA, the shrimp fed with 0.75 and 1% MG had the highest PA activity.

Discussion

The current study used a specific blend of monoglyceride (MG) commercially available as BalanGut AQ L (BASF, Germany). The product is a synergistic combination of short (propionic and butyric) to medium (caproic, caprylic, capric and lauric acid) chain-length fatty acids. The main active ingredient is Monobutyryn, which could improve epithelium integrity, host immune defense, microbiota balance and reduce liver cholesterol (Lee et al., 2021; Nguyen et al., 2019). The results of the present study showed that dietary supplementation of MG with inclusion levels of 0.75; 1.0 and 1.5% into shrimp feed could increase the final biomass, final mean weight (FMW), and weight gain (WG) as well as to increase the production efficiency by reducing the feed conversion ratio (FCR) compared to control treatment. However, as the inclusion levels of MG increases, there is a tendency for a declining trend on the growth performance of the shrimp. Interestingly, there is no difference in TGC among the dietary treatments used as a model to estimate the growth of

Table 4. Growth performance of pacific white shrimp *Penaeus vannamei* (Mean initial weight 1.06 ± 0.01 g) fed experimental diets for 60 d. Values represent the mean of six replicates. Results in the same columns with different superscript letter are significantly different ($P < 0.05$) based on analysis of variance followed by Tukey's multiple comparison test.

Diet code	Final Biomass (g)	Final Mean Weight (g)	Survival (%)	WG ¹ (%)	FCR ²	TGC ³
Control	79.07 ^b	8.54 ^b	92.50	8.18 ^b	1.30 ^a	0.0780
0.5 MG	75.25 ^b	8.65 ^b	87.50	8.30 ^b	1.29 ^a	0.0786
0.75 MG	93.91 ^a	9.63 ^a	97.50	9.22 ^a	1.15 ^b	0.0809
1.0 MG	90.53 ^a	9.41 ^a	96.25	9.02 ^a	1.18 ^b	0.0806
1.5 MG	90.15 ^a	9.38 ^a	96.25	9.01 ^a	1.18 ^b	0.0811
<i>p</i> -value	<0.0001	<0.0001	0.1012	0.0001	<0.0001	0.5189
PSE ⁴	3.5552	0.2375	3.9866	0.2355	0.0325	0.0022

Note: ¹ WG = Weight gain; ² FCR = Feed conversion ratio; ³ TGC = Thermal growth coefficient; ⁴ PSE = Pooled standard error




Figure 1. Total hemocyte count of Pacific white shrimp *Penaeus Vannamei* (10^6 cell mL^{-1}) at the end of feeding trial. Values represent the mean of five replicates (P -value = 0,457).




Figure 2. Lysozyme activity of Pacific white shrimp *Penaeus Vannamei* (U mL⁻¹) at the end of feeding trial. Values represent the mean of eight replicates (P-value = 0,055).




Figure 3. Representative histopathological images of hematoxylin and eosin-stained sections of hepatopancreas of shrimp after feeding with the experimental diets for 60-days




Figure 4. (A) Survival trend over time after challenged with *Vibrio harveyi* infection at dose of 1×10^5 CFU mL⁻¹, intramuscularly. (B) cumulative mean survival rate of shrimp at the end of challenge test. Different letters indicate statistically significant differences ($P < 0.05$) among the dietary treatment.




Figure 5. Phenoloxidase activity of experimental shrimp survivor at day-5 post injection with *Vibrio harveyi* at the dose of 1×10^5 CFU shrimp⁻¹. Different letters indicate statistically significant differences ($P < 0.05$) among the dietary treatment.

aquatic organisms in temperature range conditions. A study with short-chain fatty acid glycerides (propionate and butyrate) showed that the supplementation of these substances was able to improve the final weight and weekly weight gain of shrimp *P. vannamei* after 47 days of the cultivation period (Bruno Correa da Silva et al., 2016). Furthermore, the author highlighted that better feed efficiency, survival, and yield were obtained using 2% inclusion levels of butyrate (Bruno Correa da Silva et al., 2016). In addition, another study revealed that the inclusion of a 2% mixture of organic acids, such as propionate, butyrate, fumarate, succinate, and citrate salts, was able to improve the growth performance and feed efficiency in shrimp (Bruno Corrêa da Silva et al., 2016). However, the inclusion of 0.5% commercial products of short-and-medium fatty acid in the diet did not significantly improve the growth of gilthead sea bream *Sparus aurata* stocked with about 60 g mean initial body and cultured in 2 m³ of fiberglass tank (Rimoldi et al., 2018). Recent research in shrimp *P. vannamei* showed that after feeding for 53 days with a

diet supplemented with a mixture of glycerol esters of short-chain fatty acid (SCFAs) and medium-chain fatty acids (MCFAs), the growth of shrimp fed with 0.07 of these mixtures (SMCFAM) are lower compared to the growth of shrimp fed with 0.035 and 0.055 % of SMCFAM, and then spiked again with the inclusion of 0.11 % SMCFAM (Shin et al., 2023). In addition to the amount of MG included in the diet, further research to evaluate the inclusion effect of MG on the growth of shrimp cultured in an outdoor pond will be required to validate the results.

The total hemocytes in shrimp are vital as the first defense mechanism. In the present study, despite no statistically significant impact on the total hemocyte count (THC), we can see the increasing number of THC as the inclusion of MG increases from 0.5 to 1.0% in the diet formulation. Interesting to observe is that the trend for THC decreases as the inclusion levels go up to 1.5%. Recently, there was also no notable increase of THC in shrimp *P. vannamei* fed with several inclusion levels of propionate and butyrate from 0.5 to 2.0% for 47 days




Figure 6. Mean percentage of phagocytic activity (A) and phagocytic index (B) in shrimp blood cells after challenge test with *vibrio harveyi* at the dose of 1 x10⁵ CFU shrimp⁻¹. Different letters indicate statistically significant differences (P=0.0187) among the dietary treatment.

(Bruno Corrêa da Silva et al., 2016; Bruno Correa da Silva et al., 2016). However, studies by Bruno Corrêa da Silva et al. (2016) showed that the supplementation of butyrate and polyhydroxybutyrate as much as 2.0 % for 42 days enhances the number of THC in shrimp. Another study by Bolívar Ramírez et al. (2017) revealed that including 2.0 % of butyrate fed to shrimp for 28 days could enhance the THC in shrimp compared to the control treatment. These results show dynamics in THC numbers at different sampling points. Therefore, it is essential to have different sampling points when using a single or mixture of short- and medium-chain length fatty acids.

Lysozyme is an enzyme that is part of the innate immune system and protects shrimp from the invasion of bacterial pathogens (Hikima et al., 2003; Jollès & Jollès, 1984). Lysozyme in shrimp can be modulated; one way is by using feed additives and supplements. (Deng et al., 2013; Novriadi et al., 2021). As shown in Figure 2, despite statistical insignificance, an increasing number of lysozyme activity is observed as the inclusion levels increase from 0.5% to 1.0%. However, a corresponding decrease was observed as the inclusion of MG increased to 1.5%.

In shrimp, hepatopancreas is known to be very sensitive to dietary changes due to differences in ingredient sources and production management practices (Chen et al., 2019; Prathomya et al., 2019; Wang et al., 2016). The hepatopancreas mainly consists of branched tubules and different types of epithelial cells lining the tubules (Wu et al., 2008). In our research, all shrimp, including those within the control treatment, showed that several vacuoles appeared in the tubular epithelial cells of the hepatopancreas of the shrimp. A study by Bruno Correa da Silva et al. (2016) indicated that shrimp fed with a sodium butyrate-supplemented diet had good integrity of intestinal mucous membranes, as reflected by few vacuoles, intact and well-defined cells and intercellular cells. Furthermore, a study by Silva et al. (2016) demonstrated that including PHB and butyrate as dietary supplements did not cause any damage to shrimp's intestinal structure. In this study, including MG up to 1.5 % did not cause any morphological changes to the hepatopancreas of the shrimp.

It was reported that *Vibrio*'s have been shown to cause severe mortalities in the aquaculture industry worldwide (Novriadi, 2016; Toranzo & Barja, 1990), including in shrimp (Kewcharoen & Srisapoom, 2019; Supono et al., 2019). Among the various *Vibrio* spp., *Vibrio harveyi* represents the most frequent pathogen causing substantial production losses to the penaeid industry (Peeralil et al., 2020; Soto-Rodriguez et al., 2012). In this study, the survival trend of the shrimp over time after being challenged with *V. harveyi* intramuscularly at a dose of 1×10^5 CFU mL⁻¹ was significantly higher in the group of shrimps fed with MG compared to the control treatment. After the challenge test, we found that the phenoloxidase (PO) as well as

phagocytic activity (PA) and phagocytic index (PI) of shrimp fed with MG were significantly higher than the control treatment. Shrimp depends mostly on the innate immune systems and may have some form of adaptive immunity in defending against pathogens (Aguirre-Guzman et al., 2009; Roth & Kurtz, 2009). Subsequently, after the first defense mechanisms cannot control pathogens' access, cellular component complements with humoral responses are deployed to initiate the protective mechanisms (Kulkarni et al., 2021). Hemocytes are essential parameters involved in cellular immune responses like phagocytosis, nodule formation and encapsulation, while the prophenoloxidase (proPO) cascades and releases antimicrobial peptides and lysozyme for self-protection are two important components of humoral immune reactions in shrimp (Chen et al., 2014). The higher number of PA, PI, and PO activities post-challenge, together with the higher lysozyme activity at the end of the growth trial, indicate that MG could induce the activation of the immune system in shrimp. Study from Romano et al. (2015) indicate that the increase of PO activity with the addition of organic acid blend in diets at 10 – 20 g Kg⁻¹ of feed linearly increases the resistance of *P. vannamei* against *V. harveyi*. In addition, the inclusion of 2 g Kg⁻¹ tributyrin can significantly enhance the non-specific immune system of *P. vannamei* against *V. harveyi* during the challenge test (Lee et al., 2021).

Monoglycerides are amphipathic molecules and thus can penetrate bacterial cell membranes and alter their permeability, causing cell lysis. In addition, aquaglyceroporins in bacterial cell walls play an essential role since they enable the entry of glycerol, which is naturally used as an energy source by the bacteria. Once inside, bacterial lipases hydrolyze the ester bonds and release the fatty acid, which will then act as any other organic acid, disrupting the electron transport chain and uncoupling oxidative phosphorylation, inhibiting membrane enzymatic activities and nutrient uptake (Kabara et al., 1972; Namkung et al., 2011; Yoon et al., 2018). Therefore, it is assumed that the synergistic effect of aminobutyric, other glycerides, and glycerol in MG enhanced the non-specific immune system and the ability to disrupt the bacterial cell membrane, resulting in adequate protection and a better survival rate for shrimp against the pathogen.

The present study shows that dietary supplementation of optimized mixtures of monobutyryl, other glycerides, and glycerol (mono-, di- and triglycerides of short- and medium-chain organic acids) in relatively low inclusion levels can improve growth performance, disease resistance against *V. harveyi*, phenoloxidase activity, phagocytic index and activity as well as the health condition in shrimp. The results further show that MG could be a potential supplement in developing functional feed for shrimp. The optimum dietary level for *P. vannamei* will likely be 0.75 – 1.0% MG in the diet formulation.

Ethical Statement

All procedures were conducted in accordance with the Indonesian Animal (Scientific Procedures) SNI 01-7252-2006 and SNI 8008:2014, approved by institutional ethical review committees (Politeknik Ahli Usaha Perikanan)

Funding Information

BASF provided monoglycerides in the form of BalanGut® AQ L for the production of experimental diet as well as funding for this study.

Author Contribution

RN: Research conceptualization, Methodology, Validation, formal analysis, investigation, Draft creation, supervision and writing the original draft.

NK: Research conceptualization, Methodology, Validation, formal analysis, investigation, Draft creation, supervision and writing the original draft.

ASK: Research conceptualization, Methodology, Validation, formal analysis, investigation, Draft creation, supervision and writing the original draft.

VEH: Research conceptualization, Methodology, Validation, formal analysis, investigation, Draft creation, supervision and writing the original draft.

SBP: Research conceptualization, Methodology, Validation, formal analysis, investigation, Draft creation, and visualization.

SW: Research conceptualization, Methodology, Validation, formal analysis, investigation, Draft creation, and visualization.

CM: Research conceptualization, Methodology, Validation, formal analysis, investigation, and funding acquisition.

AB: Research conceptualization and funding acquisition.

PS: Research conceptualization and funding acquisition.

Conflict of Interest

Dr. Chi Man and Peer Staehler was employed by BASF and Dr. Adriana Barri still under BASF. The remaining authors state no conflict of interest.

Acknowledgements

The work was supported by the grants from BASF (Germany). Special thanks to the students from the Faculty of Fisheries and Marine Sciences, Diponegoro University, Semarang, Indonesia, for their help in preparing the experimental tanks. The authors would also like to extend their gratitude to those who have taken the time to critically review this manuscript and those who helped support this research. Mentioning a trademark or proprietary product does not constitute an endorsement. It does not imply its approval to the exclusion of other products that may also be suitable.

References

- Abo-Taleb, H. A., El-feky, M. M., Azab, A. M., Mabrouk, M. M., Elokaby, M. A., Ashour, M., Mansour, A. T., Abdelzaher, O. F., Abualnaja, K. M., & Sallam, A. E. (2021). Growth performance, feed utilization, gut integrity, and economic revenue of grey mullet, *Mugil cephalus*, fed an increasing level of dried zooplankton biomass meal as fishmeal substitutions. *Fishes*, *6*(3), 38.
- Aguirre-Guzman, G., Sanchez-Martinez, J. G., Campa-Cordova, A. I., Luna-Gonzalez, A., & Ascencio, F. (2009). Penaeid shrimp immune system. *The Thai Journal of Veterinary Medicine*, *39*(3), 205-215.
- Al Eissa, A., Chen, P., Brown, P. B., & Huang, J. Y. (2022). Effects of feed formula and farming system on the environmental performance of shrimp production chain from a life cycle perspective. *Journal of Industrial Ecology*, *26*(6), 2006-2019.
- Amend, D. F. (1981). Potency testing of fish vaccines. *Fish biologics: serodiagnostics and vaccines*, 447-454.
- Ayiku, S., Shen, J.-f., Tan, B.-p., Dong, X.-h., & Liu, H.-y. (2020). Effects of dietary yeast culture on shrimp growth, immune response, intestinal health and disease resistance against *Vibrio harveyi*. *Fish & Shellfish Immunology*, *102*, 286-295.
- Bolívar Ramírez, N. C., Rodrigues, M. S., Guimarães, A. M., Guertler, C., Rosa, J. R., Seiffert, W. Q., Andreatta, E. R., & Vieira, F. d. N. (2017). Effect of dietary supplementation with butyrate and probiotic on the survival of Pacific white shrimp after challenge with *Vibrio alginolyticus*. *Revista Brasileira de Zootecnia*, *46*, 471-477.
- Chen, S., Zhuang, Z., Yin, P., Chen, X., Zhang, Y., Tian, L., Niu, J., & Liu, Y. (2019). Changes in growth performance, haematological parameters, hepatopancreas histopathology and antioxidant status of pacific white shrimp (*Litopenaeus vannamei*) fed oxidized fish oil: Regulation by dietary myo-inositol. *Fish & Shellfish Immunology*, *88*, 53-64.
- Chen, Y.-Y., Chen, J.-C., Lin, Y.-C., Putra, D. F., Kitikiew, S., Li, C.-C., Hsieh, J.-F., Liou, C.-H., & Yeh, S.-T. (2014). Shrimp that have received carrageenan via immersion and diet exhibit immunocompetence in phagocytosis despite a post-plateau in immune parameters. *Fish & Shellfish Immunology*, *36*(2), 352-366.
- da Silva, B. C., Jatobá, A., Schleder, D. D., Vieira, F. d. N., Mourinho, J. L. P., & Seiffert, W. Q. (2016). Dietary supplementation with butyrate and polyhydroxybutyrate on the performance of pacific white shrimp in biofloc systems. *Journal of the World Aquaculture Society*, *47*(4), 508-518.
- da Silva, B. C., Vieira, F. d. N., Mourinho, J. L. P., Bolivar, N., & Seiffert, W. Q. (2016). Butyrate and propionate improve the growth performance of *Litopenaeus vannamei*. *Aquaculture Research*, *47*(2), 612-623.
- Delphino, M. K., Laurin, E., Patanasatienkul, T., Rahardjo, R. B., Hakim, L., Zulfikar, W. G., Burnley, H., Hammell, K. L., & Thakur, K. (2022). Description of biosecurity practices on shrimp farms in Java, Lampung, and Banyuwangi, Indonesia. *Aquaculture*, *556*, 738277.
- Deng, D., Mei, C., Mai, K., Tan, B. P., Ai, Q., & Ma, H. (2013). Effects of a yeast-based additive on growth and immune responses of white shrimp, *Litopenaeus vannamei* (Boone, 1931), and aquaculture environment. *Aquaculture Research*, *44*(9), 1348-1357.

- Fox, J., Treece, G., & Sanchez, D. (2001). Shrimp nutrition and feed management. *Methods for improving shrimp farming in Central America*, 65-90.
- Fox, J. M., Lawrence, A. L., & Smith, F. (2004). Development of a low-fish meal feed formulation for commercial production of *Litopenaeus vannamei*. *Avances en nutrición acuicola*.
- Gatlin III, D. M., Barrows, F. T., Brown, P., Dabrowski, K., Gaylord, T. G., Hardy, R. W., Herman, E., Hu, G., Krogdahl, Å., & Nelson, R. (2007). Expanding the utilization of sustainable plant products in aquafeeds: a review. *Aquaculture Research*, 38(6), 551-579.
- Geetha, R., Avunje, S., Solanki, H., Priyadarshini, R., Vinoth, S., Anand, P., Ravisankar, T., & Patil, P. K. (2022). Farm-level economic cost of Enterocytozoon hepatopenaei (EHP) to Indian *Penaeus vannamei* shrimp farming. *Aquaculture*, 548, 737685.
- Gyan, W. R., Yang, Q.-h., Tan, B., Xiaohui, D., Chi, S., Liu, H., & Zhang, S. (2021). Effects of replacing fishmeal with dietary dried distillers grains with solubles on growth, serum biochemical indices, antioxidative functions, and disease resistance for *Litopenaeus vannamei* juveniles. *Aquaculture Reports*, 21, 100821.
- Hikima, S., Hikima, J.-i., Rojtinnakorn, J., Hirono, I., & Aoki, T. (2003). Characterization and function of kuruma shrimp lysozyme possessing lytic activity against *Vibrio* species. *Gene*, 316, 187-195.
- Huang, J., Yang, Y., & Wang, A. (2010). Reconsideration of phenoloxidase activity determination in white shrimp *Litopenaeus vannamei*. *Fish & Shellfish Immunology*, 28(1), 240-244.
- Jin, M., Xiong, J., Zhou, Q.-C., Yuan, Y., Wang, X.-X., & Sun, P. (2018). Dietary yeast hydrolysate and brewer's yeast supplementation could enhance growth performance, innate immunity capacity and ammonia nitrogen stress resistance ability of Pacific white shrimp (*Litopenaeus vannamei*). *Fish & Shellfish Immunology*, 82, 121-129.
- Jithendran, K. P., Navaneeth Krishnan, A., Jagadeesan, V., Anandaraja, R., Ezhil Praveena, P., Anushya, S., Bala Amarnath, C., & Bhuvaneshwari, T. (2021). Co-infection of infectious myonecrosis virus and Enterocytozoon hepatopenaei in *Penaeus vannamei* farms in the east coast of India. *Aquaculture Research*, 52(10), 4701-4710.
- Jollès, P., & Jollès, J. (1984). What's new in lysozyme research? Always a model system, today as yesterday. *Molecular and cellular biochemistry*, 63, 165-189.
- Kabara, J. J., Swieczkowski, D. M., Conley, A. J., & Truant, J. P. (1972). Fatty acids and derivatives as antimicrobial agents. *Antimicrobial agents and chemotherapy*, 2(1), 23-28.
- Kewcharoen, W., & Srisapoome, P. (2019). Probiotic effects of *Bacillus* spp. from Pacific white shrimp (*Litopenaeus vannamei*) on water quality and shrimp growth, immune responses, and resistance to *Vibrio parahaemolyticus* (AHPND strains). *Fish & shellfish immunology*, 94, 175-189.
- Kok, B., Malcorps, W., Tlustý, M. F., Eltholth, M. M., Auchterlonie, N. A., Little, D. C., Harmsen, R., Newton, R. W., & Davies, S. J. (2020). Fish as feed: Using economic allocation to quantify the Fish In: Fish Out ratio of major fed aquaculture species. *Aquaculture*, 528, 735474.
- Kulkarni, A., Krishnan, S., Anand, D., Kakkattunivarthil Uthaman, S., Otta, S. K., Karunasagar, I., & Kooloth Valappil, R. (2021). Immune responses and immunoprotection in crustaceans with special reference to shrimp. *Reviews in Aquaculture*, 13(1), 431-459.
- Lee, C., Shin, J., Feyaerts, J., Shin, J., Kim, M. G., Gunathilaka, B. E., Ko, D., Jo, S., Eom, G., & Ji, S. C. (2021). Effects of dietary supplementation of monobutyryl and tributyrin on growth, feed efficiency, innate immunity, digestibility and disease resistance of Pacific white shrimp (*Litopenaeus vannamei*) against *Vibrio harveyi*. *Aquaculture nutrition*, 27(3), 771-781.
- Lim, C., & Dominy, W. (1990). Evaluation of soybean meal as a replacement for marine animal protein in diets for shrimp (*Penaeus vannamei*). *Aquaculture*, 87(1), 53-63.
- López-Téllez, N. A., Corbalá-Bermejo, J. A., Bustamante-Unzueta, M. L., Silva-Ledesma, L. P., Vidal-Martínez, V. M., & Rodríguez-Canul, R. (2020). History, impact, and status of infectious diseases of the Pacific white shrimp *Penaeus vannamei* (Bonne, 1831) cultivated in Mexico. *Journal of the World Aquaculture Society*, 51(2), 334-345.
- Miles, R. D., & Chapman, F. A. (2006). The benefits of fish meal in aquaculture diets. *EDIS*, 2006(12).
- Namkung, H., Yu, H., Gong, J., & Leeson, S. (2011). Antimicrobial activity of butyrate glycerides toward *Salmonella Typhimurium* and *Clostridium perfringens*. *Poultry science*, 90(10), 2217-2222.
- Nguyen, T. D., Prykhodko, O., Hållénus, F. F., & Nyman, M. (2019). Monobutyryl reduces liver cholesterol and improves intestinal barrier function in rats fed high-fat diets. *Nutrients*, 11(2), 308.
- Nguyen, T. N., Davis, D. A., & Saoud, I. P. (2009). Evaluation of alternative protein sources to replace fish meal in practical diets for juvenile tilapia, *Oreochromis* spp. *Journal of the World Aquaculture Society*, 40(1), 113-121.
- Niu, J., Xie, J.-J., Guo, T.-Y., Fang, H.-H., Zhang, Y.-M., Liao, S.-Y., Xie, S.-W., Liu, Y.-J., & Tian, L.-X. (2019). Comparison and evaluation of four species of macro-algae as dietary ingredients in *Litopenaeus vannamei* under normal rearing and WSSV challenge conditions: effect on growth, immune response, and intestinal microbiota. *Frontiers in physiology*, 9, 1880.
- Novriadi, R. (2016). Vibriosis in aquaculture. *Omni-Akuatika*, 12(1).
- Novriadi, R., Fadhillah, R., Wahyudi, A. E., & Trullas, C. (2021). Effects of hydrolysable tannins on the growth performance, total haemocyte counts and lysozyme activity of pacific white leg shrimp *Litopenaeus vannamei*. *Aquaculture Reports*, 21, 100796.
- Novriadi, R., Istiqomah, I., Isnansetyo, A., Balk, D., Jolly-Breithaupt, M., & Davies, S. (2023). Corn fermented protein in production diets for pacific white legged shrimp *Litopenaeus vannamei*: Improved growth performance, health status and resistance to infection. *Aquaculture Reports*, 30, 101571.
- Novriadi, R., Roigé, O., & Segarra, S. (2022). Effects of dietary nucleotide supplementation on performance, profitability, and disease resistance of *Litopenaeus vannamei* cultured in Indonesia under intensive outdoor pond conditions. *Animals*, 12(16), 2036.
- Novriadi, R., Salze, G., Abebe, A., Hanson, T., & Davis, D. A. (2019). Partial or total replacement of fish meal in the diets of Florida pompano *Trachinotus carolinus*. *Aquaculture Research*, 50(5), 1527-1538.
- Peeralil, S., Joseph, T. C., Murugadas, V., Akhila, P., Sreejith, V., & Lalitha, K. (2020). *Vibrio harveyi* virulence gene expression in vitro and in vivo during infection in black

- tiger shrimp *Penaeus monodon*. *Diseases of aquatic organisms*, 139, 153-160.
- Prathomya, P., Hussein, H. A., Tran, N. T., Duan, Y. H., Peng, S. H., Xu, Y., Qian, X. Q., & Wang, G. T. (2019). Effects of dietary supplementation of *Astragalus membranaceus*, *Codonopsis pilosula*, and *Glycyrrhiza uralensis* extract mixture on growth performance, haematological parameters and hepatopancreatic performance in juvenile Pacific white shrimp (*Litopenaeus vannamei*). *Aquaculture Research*, 50(9), 2707-2717.
- Rahman, R. A., Kendall, G., Ramli, R., Jamari, Z., & Ku-Mahamud, K. R. (2017). Shrimp feed formulation via evolutionary algorithm with power heuristics for handling constraints. *Complexity*, 2017.
- Rimoldi, S., Gliozheni, E., Ascione, C., Gini, E., & Terova, G. (2018). Effect of a specific composition of short-and medium-chain fatty acid 1-Monoglycerides on growth performances and gut microbiota of gilthead sea bream (*Sparus aurata*). *PeerJ*, 6, e5355.
- Romano, N., Koh, C.-B., & Ng, W.-K. (2015). Dietary microencapsulated organic acids blend enhances growth, phosphorus utilization, immune response, hepatopancreatic integrity and resistance against *Vibrio harveyi* in white shrimp, *Litopenaeus vannamei*. *Aquaculture*, 435, 228-236.
- Roth, O., & Kurtz, J. (2009). Phagocytosis mediates specificity in the immune defence of an invertebrate, the woodlouse *Porcellio scaber* (Crustacea: Isopoda). *Developmental & Comparative Immunology*, 33(11), 1151-1155.
- Shin, J., Feyaerts, J., Hasanathi, M., Lee, C., Shin, D., Lee, Y., Lim, J., Kim, H.-S., & Lee, K.-J. (2023). Evaluation of a mixture of short-chain and medium-chain fatty acid glycerides as a dietary supplement in diets for Pacific white shrimp (*Litopenaeus vannamei*). *Aquaculture International*, 31(3), 1483-1498.
- Shinn, A., Pratoomyot, J., Griffiths, D., Trong, T., Vu, N. T., Jiravanichpaisal, P., & Briggs, M. (2018). Asian shrimp production and the economic costs of disease. *Asian Fish. Sci. S*, 31, 29-58.
- Silva, B., Jesus, G., Seiffert, W., Vieira, F., Mourinho, J., Jatobá, A., & Nolasco-Soria, H. (2016). The effects of dietary supplementation with butyrate and polyhydroxybutyrate on the digestive capacity and intestinal morphology of Pacific White Shrimp (*Litopenaeus vannamei*). *Marine and Freshwater Behaviour and Physiology*, 49(6), 447-458.
- Soto-Rodriguez, S. A., Gomez-Gil, B., Lozano, R., del Rio-Rodríguez, R., Diéguez, A. L., & Romalde, J. L. (2012). Virulence of *Vibrio harveyi* responsible for the "Bright-red" Syndrome in the Pacific white shrimp *Litopenaeus vannamei*. *Journal of invertebrate pathology*, 109(3), 307-317.
- Supono, S., Wardiyanto, W., & Harpeni, E. (2019). Identification of *Vibrio* sp. as a cause of white feces diseases in white shrimp *Penaeus vannamei* and handling with herbal ingredients in East Lampung Regency, Indonesia. *AAFL Bioflux*, 12(2), 417-425.
- Tacon, A. G. (1993). Feed ingredients for warmwater fish, fish meal and other processed feedstuffs. *FAO Fisheries Circular (FAO)*. no. 856.
- Toranzo, A. E., & Barja, J. L. (1990). A review of the taxonomy and seroepizootiology of *Vibrio anguillarum*, with special reference to aquaculture in the northwest of Spain. *Diseases of aquatic organisms*, 9(1), 73-82.
- Wang, X., Li, E., Xu, C., Qin, J. G., Wang, S., Chen, X., Cai, Y., Chen, K., Gan, L., & Yu, N. (2016). Growth, body composition, ammonia tolerance and hepatopancreas histology of white shrimp *Litopenaeus vannamei* fed diets containing different carbohydrate sources at low salinity. *Aquaculture Research*, 47(6), 1932-1943.
- Wu, J.-P., Chen, H.-C., & Huang, D.-J. (2008). Histopathological and biochemical evidence of hepatopancreatic toxicity caused by cadmium and zinc in the white shrimp, *Litopenaeus vannamei*. *Chemosphere*, 73(7), 1019-1026.
- Xie, J. J., Lemme, A., He, J. Y., Yin, P., Figueiredo-Silva, C., Liu, Y. J., Xie, S. W., Niu, J., & Tian, L. X. (2018). Fishmeal levels can be successfully reduced in white shrimp (*Litopenaeus vannamei*) if supplemented with DL-Methionine (DL-Met) or DL-Methionyl-DL-Methionine (Met-Met). *Aquaculture nutrition*, 24(3), 1144-1152.
- Yoon, B. K., Jackman, J. A., Valle-González, E. R., & Cho, N.-J. (2018). Antibacterial free fatty acids and monoglycerides: biological activities, experimental testing, and therapeutic applications. *International journal of molecular sciences*, 19(4), 1114.
- Zinn, K. E., Hernot, D., Fastinger, N., Karr-Lilienthal, L., Bechtel, P., Swanson, K., & Fahey Jr, G. (2009). Fish protein substrates can substitute effectively for poultry by-product meal when incorporated in high-quality senior dog diets. *Journal of animal physiology and animal nutrition*, 93(4), 447-455.